
The origin of σ -drops: mapping stellar kinematics and populations in spirals

Eric Emsellem¹

Université de Lyon 1, CRAL, Observatoire de Lyon, 9 av. Charles André, F-69230 Saint-Genis Laval; CNRS, UMR 5574 ; ENS de Lyon, France
 emsellem@obs.univ-lyon1.fr

1 Introduction

The link between nuclear activity and the host galaxy remains elusive. It seems now clear that the galactic environment is not the key, and that there is no significant apparent difference in terms of the presence of bars/spirals between active and non-active galaxies. Nuclear activity nevertheless requires a small central gas reservoir (10^4 - 10^7 $M_\odot$) which may be only partly consumed during the $\sim 10^6 - 10^7$ yr AGN duty cycle. A possible route towards a better understanding of the involved physical processes is to directly probe the gravitational potential of the host galaxies, with the hope that the corresponding sensitive tracers (stars, gas) will deliver convincing evidence for a kinematic signature associated with the AGN. This is the motivation for an on-going survey of 50 galaxies, for which the two-dimensional kinematics of the (neutral, ionised and molecular) gas and stellar components are being obtained (see Dumas et al., this conference). With only 10 to 15% of Seyfert galaxies in the local Universe, we must also acknowledge the possibility that this activity is a recurrent (but short) process, and that signatures of recent gas accretion associated with the onset of the central activity should be detectable. This paper is a short report on the detection and study of σ -drops which we believe are the result of past gas accretion followed by subsequent star formation.

2 σ -drops: the list is growing

In the course of an observational program aimed at revealing dynamical evidences for the presence of inner bars in spiral galaxies, we have detected a decrease of the stellar velocity dispersion towards the centre of 3 out of 3 barred galaxies [2] where we could probe the very central region. We suggested that these σ -drops are the result of gas accretion followed by star formation, and therefore indicative of a relatively recent infall of dissipative material.

This idea was successfully tested for the specific case of an infall driven by a large-scale bar [7], although this does not mean that bars are the preferred cause for such a process.

Since this study of a very small sample of barred galaxies, the list of σ -drops has been growing very significantly (and steadily). Fig 1 provides a non-exhaustive list of such objects and shows that σ -drops are indeed a rather common phenomenon. Most of the galaxies listed in Fig 1 contain active nuclei, but this is most certainly due to a bias in the observed sample where reliable stellar kinematics have been obtained.

NGC 1068	SBb	Sy2	Emsellem et al. 2006
NGC 1097	SB(s)b	Sy2	Emsellem et al. 2001
NGC 1138	SBO	?	Simien & Prugniel 2002
NGC 1808	SAB(s:)b	Sy 2/starburst	Emsellem et al. 2001
NGC 2639	SAa	Sy2	Marquez et al. 2003
NGC 2460	SAa	?	Shapiro et al. 2003
NGC 2775	SAab	?	Shapiro et al. 2003
NGC 3021	SAbc		Héraudeau et al. 1999
NGC 3412	SB0		Agueri et al. 2003
NGC 4030	SAbc	?	Shapiro et al. 2003
NGC 2639	SA(r)a	Sy 1.9	Marquez et al. 2003
NGC 3412	SB0(s)	?	Agueri et al. 2003
NGC 3593	SA(s)0/a	Sy 2	Bertola et al. 1993
NGC 3623	SAB(rs)a	Liner	De Zeeuw et al. 2002
NGC 3627	SBb	Liner/Sy 2	Héraudeau & Simien 1998
NGC 4303	SBbc	Sy 2	Héraudeau & Simien 1998
NGC 4579	SBb	Liner/Sy 1.9	Héraudeau & Simien 1998
NGC 4725	SBab	Sy 2	Héraudeau et al. 1999
NGC 4477	SB0	Sy 2	Jarvis et al. 1988
NGC 5728	SABb(r)	Sy 2	Emsellem et al. 2001
NGC 6503	SA(s)cd	Liner/HII	Bottema 1993; Bottema & Gerritsen 1997
NGC 6814	SAB(rs)bc	Sy 1.5	Marquez et al. 2003
NGC 6951	SAB(rs)bc	Sy 2	Marquez et al. 2003
IC 184	SB(r)a	Sy 2	Marquez et al. 2003
+ 10			Chung & Bureau 2004

Fig. 1. A (non-exhaustive) list of galaxies exhibiting a σ -drop with associated references. The columns are, from left to right: name of the galaxy, Hubble type, activity type and reference.

3 Formation scenarios

Numerical simulations including stars, gas and star formation are very helpful in constraining the formation process of σ -drops in the central regions of disc

galaxies. Recent experiments [6] clearly showed that σ -drops can be formed on a relatively short timescale (500 Myr) and are long-lived if star formation is sustained at a level of about $1 M_{\odot}/\text{yr}$ in the nuclear region. This easily occurs in a gas-rich spiral galaxy where a bar can funneled gas towards the central 500 pc or so. When star formation stops, the amplitude of the σ -drop starts to significantly decrease, as shown in Fig. 2 [6].




Fig. 2. Amplitude of the σ -drop in numerical simulations of spiral galaxies, where central star formation is sustained via regular gas infall (black curve), or stopped at different times (red, green, blue). The magenta curve corresponds to a similar simulation but where star formation is not turned on. Extracted from [6].

4 A 3D view

Integral-field spectrographs (IFS) are now routinely used in order to probe the full extent and shape of σ -drops, and hence to better understand the stellar component with which they are associated. Recent IFS data obtained on the prototypical Seyfert 2 galaxy NGC 1068 clearly demonstrated how critical is such a detailed mapping of the kinematics in the central region [1, 4]. Surveys of spiral galaxies using integral-field spectrographs [3] are now showing that about one third of spirals are expected to exhibit a σ -drop, this

being a lower limit considering the difficulty of such a detection when dust is at play. This significant fraction of galaxies with σ -drops would suggest the presence of relatively young stars associated with the underlying dynamically cold stellar component. However, there seems to be no direct relation between the presence of a σ -drop and the mean luminosity weighted age of the stellar population. In order to further test this issue, we have obtained high spectral resolution ($R = 12000$) integral-field data of a known σ -drop galaxy, NGC 3623, using FLAMES at the VLT. Fig. 3 shows the corresponding two-dimensional stellar kinematics for this galaxy, emphasizing the presence of the central σ -drop in this galaxy. A preliminary analysis of the stellar populations in that region shows that stars associated with the σ -drop cannot be younger than 1 Gyr, confirming the fact that σ -drops can be long-lived.




Fig. 3. VLT/FLAMES maps of the galaxy NGC 3623. From left to right: reconstructed image, stellar velocity field and stellar velocity dispersion field. The σ -drop is obvious and associated with the pinching of the (disk-like) isoveLOCITIES.

5 Conclusions

σ -drops are ubiquitous in spirals, with a relative fraction of about 30% or more. This suggests that significant episodes of star formation occur recurrently in the central part of these galaxies. This could in fact be linked with the common presence of blue nuclei in late-type galaxies [5]. The detection of a young stellar component associated with a σ -drop is rather difficult and requires high spectral resolution and high signal-to-noise ratio data. We have recently started a more systematic study of these stellar populations using the near-infrared integral-field spectrograph SINFONI. The analysis of these data is on-going, and should help us to constrain the star formation history in the central few hundred parsecs. This may eventually lead to a better understanding of the chronology/synchronicity between star formation and activity in the nuclear region.

References

1. Emsellem, E., Fathi, K., Wozniak, H., Ferruit, P., Mundell, C. G., & Schinnerer, E.: MNRAS **365**, 367 (2006)
2. Emsellem, E., Greusard, D., Combes, F., Friedli, D., Leon, S., Pécontal, E., & Wozniak, H.: A&A **368**, 52 (2001)
3. Ganda, K., Falcón-Barroso, J., Peletier, R. F., Cappellari, M., Emsellem, E., McDermid, R. M., de Zeeuw, P. T., & Carollo, C. M.: MNRAS **367**, 46 (2006)
4. Gerssen, J., Allington-Smith, J., Miller, B. W., Turner, J. E. H., & Walker, A.: MNRAS **365**, 29 (2006)
5. Rossa, J., van der Marel, R. P., Böker, T., Gerssen, J., Ho, L. C., Rix, H.-W., Shields, J. C., & Walcher, C.-J.: AJ **132**, 1074 (2006)
6. Wozniak, H., & Champavert, N.: MNRAS **369**, 853 (2006)
7. Wozniak, H., Combes, F., Emsellem, E., & Friedli, D.: A&A **409**, 469 (2003)